



(21) (A1) **2,228,049**  
(86) 1996/07/29  
(87) 1997/02/13

(72) SCHLACHTER, Herbert, DE

(72) HAMM, Michael, DE

(71) SCHLACHTER, Herbert, DE

(71) HAMM, Michael, DE

(51) Int.Cl.<sup>6</sup> A61K 35/78, A61K 33/06, A23L 1/29, A61K 31/68, A61K 31/20

(30) 1995/07/28 (95111978.3) EP

(54) **PREPARATION A DEUX PHASES**

(54) **TWO-PHASE PREPARATION**

(57) La présente invention concerne une préparation à deux phases, destinée à un emploi échelonné dans le temps, qui comprend un produit A pour la première phase et un produit B pour la deuxième phase. Ces produits comportent chacun, indépendamment l'un de l'autre, au moins un acide gras non saturé et/ou au moins un élément du groupe des oligo-éléments et des substances minérales et/ou au moins un élément du groupe des vitamines et/ou au moins un élément du groupe des substances végétales biologiquement actives, par exemple des polyphénols, des bioflavonoïdes ou des fibres alimentaires et/ou au moins un acide aminé et/ou un dérivé d'acide aminé. Le produit A et/ou le produit B peuvent comporter en outre, le cas échéant, de la lécithine de soja, A et B devant toutefois se distinguer par leur composition quantitative ou par la nature de leurs constituants. La préparation de l'invention peut être employée comme complément nutritif (produit diététique) ou comme médicament.

(57) The invention concerns a preparation designed to act gradually and containing a product A for the first stage and a product B for the second stage. Each of the products includes, independently of each other, at least one unsaturated fatty acid and/or at least one trace element or mineral and/or at least one vitamin and/or at least one bioactive plant substance, e.g. a polyphenol, biflavonoid or dietary fibre, and/or at least one amino acid and/or amino-acid derivative, product A and/or product B optionally also containing soya lecithin, with the provision that product A and product B differ from each other in their amounts and/or their material composition. The preparation can be used as a dietary complement or as a drug.



PCT/EP96/03329  
Schlachter, Herbert, et al.  
u.Z.: A 2376 PCT

#### Two-Phase Preparation

The invention relates to a two-phase preparation for use at differential times containing a product A for the first phase and a product B for the second phase which comprise, independently of each other, at least one representative of the unsaturated fatty acids and/or at least one representative of the group of trace elements and minerals and/or at least one representative of the group of vitamins and/or at least one representative of the group of bioactive plant substances, e.g. polyphenoles, bioflavonoids or dietary fiber, and/or at least one amino acid and/or amino acid derivative, product A and/or product B optionally additionally containing soy lecithin, with the proviso that product A and product B differ from each other in their quantitative and/or their material composition. The preparation can be used as a food supplement (dietary supplement) or as a drug.

In particular, the invention relates to a two-phase preparation for use at differential times comprising a product A and a product B which include, independently of each other, at least one representative of the unsaturated fatty acids, one representative of the group of trace elements and minerals and one representative of the group of

vitamins, product A and/or product B optionally additionally containing soy lecithin, and/or at least one representative of the group of bioactive plant substances and/or at least one amino acid and/or amino acid derivative which is used as a food supplement (dietary supplement).

It is generally known that the nutrients carbohydrates as well as the dietary fats are mainly used to cover the body's energy requirement. Proteins are important building components for cells and endogenous active substances such as enzymes and certain hormones. Polyunsaturated fatty acids, vitamins, minerals and trace elements as well as bioactive plant substances such as flavonoids are becoming more and more important for health and fitness.

Vitamins, minerals and trace elements, polyunsaturated fatty acids and bioactive plant substances such as flavonoids are essential regulators in the metabolism and nutrients protecting health. However, a one-sided diet, stringent weight reduction diets, preference of high-calorie foods of low nutrient density and the high degree of processing of many foods but also the partial impoverishment of the soils with respect to important minerals make an optimal supply with all the necessary nutrients difficult. In addition, due to adverse environmental effects, smoking, consuming alcohol and medicaments as well as stress, there is an increased demand of some essential nutrients, for example magnesium, zinc, vitamin E, vitamin C, B-type vitamins and beta carotene. A food complementary nutrient substitution therefore contributes to a sufficient supply of nutrients, an optimal protection of health and well-being, as does a well-balanced nutrition which is high-roughage and low-fat. The reference "Empfehlungen für die Nährstoffzufuhr" [Recommendations for the supply with nutrients], Deutsche Gesellschaft für Ernährung, 5th edition, 1991, Umschau Verlag, Frankfurt/Main, describes recommendations for the body's supply with nutrients.

Vitamins are essential nutritional components which are more or less indispensable for the normal body functions of heterotroph organisms and which have to be supplied as required since they are only available either from outside sources or under the influence of milieu factors (e.g. intestinal flora). Their specific biocatalytic effect is to replace the active component of enzymes which is subject to metabolical consumption. From science it is for example known that B-type vitamins participate in the intermediary metabolism as coenzymes and that vitamins C, E and beta carotene mainly function as antioxidants. A deficiency due to an insufficient supply or resorption, a disorder of the intestinal flora or the metabolism, the effects of antivitamins or an increased consumption lead to hypovitaminosis and avitaminosis.

Furthermore, minerals and trace elements are essential regulators in the metabolism. Calcium is the most important mineral building substance for bones and teeth. A sufficient supply of calcium up until the age of 30 is the best protection against osteoporosis during old age. After that, sufficient food calcium ensures that the depot in the bones is not unduly strained. For women calcium is often a critical nutrient. Zinc, magnesium and B-type vitamins are high-performance elements which activate enzymes and thus allow the metabolism of carbohydrates, fats and proteins. Silicon favorably contributes to the stability and maintenance of skin, hair and nails. Furthermore, it is scientifically undisputed that zinc fulfills an essential function in the body's immune system and the skin metabolism.

Polyunsaturated fatty acids from purely vegetal sources such as for example evening primrose oil and linseed oil, contain linolic acid and  $\alpha$ - and  $\gamma$ -linolenic acid, which are important starting substances for biologically active

regulators, as for example eicosanoids and prostaglandins, and ensure a healthy equilibrium in the metabolism. Eicosanoids and prostaglandins, also referred to as tissue hormones, are currently being studied extensively by scientists for their health stabilizing effects. The favorable effect of polyunsaturated fatty acids on the cholesterol metabolism and healthy skin function as well as on inflammatory processes is known. It is furthermore known that polyunsaturated fatty acids of the type of  $\omega$ -3 (icosapentenoic acid,  $\alpha$ -linolenic acid) and  $\omega$ -6 (linolic acid,  $\gamma$ -linolenic acid) fatty acids have a favorable effect on migraine, rheumatism, neurodermatitis, psoriasis, premenstrual syndrome and regeneration processes induced by physical strain during exercise (cf. A. Berg, D. König, H. Schlachter, J. Keul, *Deutsche Zeitschrift für Sportmedizin* [German Magazine of Sports Medicine], Volume 44 (1993), Special Issue, and literature cited therein).

Carrot, broccoli and citrus extracts are natural sources of vitamins and minerals and contain the so-called secondary plant substances, which are also referred to as bioactive plant substances in the special branch of science whose significance has been recognized and is now being examined as well. These natural plant substances also include bioflavonoids which effectively support the impact of vitamin C with respect to the power of resistance, the vascular walls and the connective tissue. It is also known that bioflavonoids have antioxidant properties and therefore synergistically complement the effect of vitamins C, E and beta carotene. Green tea as well is an excellent source of these vegetal antioxidants (e.g. polyphenols) (cf. B. Watzl, C. Leitzmann *Bioaktive Substanzen in Lebensmitteln* [Bioactive substances in foodstuffs], Hippokrates Publishing House, Stuttgart, 1995).

In addition to their function as building blocks for proteins, amino acids are precursors of biologically

effective compounds, for example serotonin, growth and stress hormones. They are also said to have anticatabolic properties (cf. in this connection K.R. Geiß, M. Hamm *Handbuch Sporternährung* [Handbook of sports nutrition], Rowohlt Publishing House, 1992).

A great number of food supplements are commercially available which contain individual vitamins (e.g. vitamin E preparations) or combinations thereof (e.g. multivitamin preparations) and/or individual minerals and trace elements (e.g. magnesium or iron preparations) or combinations thereof (e.g. multimineral capsules). There are also preparations with polyunsaturated fatty acids alone or in combination with vitamins and minerals. For example, a preparation may contain many different vitamins and some selected minerals which cover the daily requirement of these nutrients.

Another commercially available preparation is characterized as a supplementary balanced diet for covering a specific increased demand of essential nutrients, in particular in view of a purpose-specific nutrition, for example after a heart attack or by-pass surgery, when the strained situation of the metabolism requires compensation. As a daily requirement, this so-called dietary foodstuff consists of a bag with 9 tablets or capsules whereby the contents of the bag are usually taken with or after a meal, whereby, however, it is also possible to take the individual capsules or tablets with meals in any desired order over the whole day. The daily requirement of this preparation comprises essential vitamins such as vitamin A, B1, B2, B3, B6, B12, C, E, E3, K and beta carotene, trace elements such as iron, zinc, manganese, chromium, magnesium and selenium, and specific essential fatty acids. However, this dietary foodstuff contains amounts of vitamins larger than the basic recommended daily allowance for a healthy adult.

Most of the commercially available food supplements are presented in the form of capsules or tablets and are usually taken independently of the time of day.

So far, food supplements have generally been intended to cover the nutritional requirements resulting from an insufficient nutrition. Despite the large number of food supplements, no optimal solution for an appropriate supply has yet been provided which sufficiently takes into account the scientific findings. From science, it is known that in contrast to a combination of many vitamins and minerals in one single form of administration, spreading the administration of these substances makes more sense from a physiological point of view. Compared to a high dosage of nutrients, the lower individual doses are more readily available. It is also known that the simultaneous intake of minerals and trace elements in a single form of administration can adversely affect the resorption of the individual components. Amongst others, magnesium and calcium as well as iron, zinc and selenium adversely affect each other.

In many known preparations, the composition of the ingredients does not always make sense as the simultaneous administration of all essential nutrients can reduce the availability of the individual components (for example, magnesium is not absorbed as well if the calcium content of a single dose is markedly higher than the magnesium content). Also, due to the prooxidative properties of iron, undesired reactions can take place between iron and polyunsaturated fatty acids.

In one of the above-mentioned commercially available preparations, the user has to take nine capsules or tablets a day with one or more meals. However, an acceptable daily dose should not exceed three individual doses each day (e.g. one to three capsules a day).

The human metabolism goes through day and night phases. It is assumed that these day and night phases also influence the nutrient balance, nutrient absorption and requirements of the body. It is an object of the present invention to provide a reasonably composed two-phase preparation wherein the daily supply of nutrients is distributed with respect to activating nutritional principles to the first half of the day and with respect to regenerative nutritional principles to the second half of the day. Another object of the present invention is to provide a two-phase preparation tailored to the needs of specific target groups. The two-phase preparation is to be used as a dietary supplement and/or a drug.

One of the objects of the present invention is achieved by providing a two-phase preparation for use at differential times comprising a product A for the first phase (active phase) and a product B for the second phase (regenerative phase) wherein the product A comprises

- a) at least one representative of the unsaturated fatty acids and/or
- b) at least one representative of the group of trace elements and minerals and/or
- c) at least one representative of the group of vitamins and/or
- d) at least one representative of the bioactive plant substances and/or
- e) at least one amino acid and/or amino acid derivative

and the product B comprises

- a) at least one representative of the unsaturated fatty acids and/or
- b) at least one representative of the group of trace elements and minerals and/or



- c) at least one representative of the group of vitamins and/or
- d) at least one representative of the group of bioactive plant substances and/or
- e) at least one amino acid and/or amino acid derivative

with the proviso that product A and product B differ from each other in their quantitative and/or material composition.

In a preferred embodiment, the present invention also provides a two-phase preparation for use at differential times comprising a product A for the first phase and a product B for the second phase wherein the product A comprises

- a) at least one representative of the unsaturated fatty acids,
- b) at least one representative of the group of trace elements and minerals and
- c) at least one representative of the group of vitamins

and the product B comprises

- a) at least one representative of the unsaturated fatty acids,
- b) at least one representative of the group of trace elements and minerals and
- c) at least one representative of the group of vitamins.

In an especially preferred embodiment, a two-phase dietary supplement for use at differential times is provided comprising a product A for the first phase and a product B for the second phase wherein the product A and the product B each comprise, independently of each other,

- a) at least one representative of the unsaturated fatty acids,
- b) at least one representative of the group of trace elements and minerals and
- c) at least one representative of the group of vitamins.

In another preferred embodiment, the product A and/or the product B can additionally comprise soy lecithin. Soy lecithin is preferably added in an amount of 1 to 40% by weight, and most preferably 5 to 20% by weight, based on the sum of all active ingredients of product A and B, respectively.

In a further embodiment, the product A and/or the product B of the preferred embodiments can additionally comprise at least one representative of the group of bioactive plant substances.

In a further embodiment, the product A and/or the product B of the preferred embodiments can additionally comprise amino acids and/or amino acid derivatives, also in combination with other active ingredients of the present invention.

The term "for use at differential times" means that the product A is to be applied or to provide its effect at the beginning of the day phase or active phase and the product B at the beginning of the night phase or regenerative phase of the human metabolism. Preferably, the product A is administered in the morning and the product B in the evening of the same day.

The term "two-phase" refers to the day or active phase as first phase (first half of the day) and to the night or regenerative phase as the second phase (second half of the day) of the human metabolism.

Product A and product B of the present invention differ from each other in their quantitative and/or material composition.

The formulations of the present invention preferably take into account synergistic effects, e.g. in the case of vitamin C and bioflavonoids as well as vitamin C and polyunsaturated fatty acids.

In the two-phase preparation of the present invention the products A and B can comprise all known representatives of the vitamins. In particular, the product A of the two-phase preparation preferably comprises the vitamins C and E as antioxidants (cell protection vitamins), and the vitamins B1, B2, B6, B12, pantothenic acid, beta carotene and niacin, and the product B preferably comprises the vitamins B1, B2, B6, B12, pantothenic acid, niacin and folic acid and biotin as metabolism regulators in the sense of coenzymes of the carbohydrate, fat and protein metabolism.

In the preferred embodiments of the two-phase preparation of the present invention, the content of vitamins is preferably 1 to 70% by weight in product A and 0.5 to 50% by weight in product B, based on the sum of all active ingredients of product A and B, respectively. In a more preferred embodiment, product A comprises 5 to 50% by weight of vitamins, especially preferred 20 to 45% by weight, and product B comprises 1 to 40% by weight of vitamins, especially preferred 1 to 35% by weight. The vitamins are added both in natural form as extracts (e.g. D- $\alpha$ -tocopherol) and in synthetic form (e.g. B-type vitamins), which, however, does not influence the effectivity of the vitamins.

Representatives of the group of trace elements and minerals in the two-phase preparation of the present invention comprises all known minerals and trace elements. In a preferred embodiment, the product A of the two-phase

preparation in particular comprises zinc, magnesium and chromium and the product B comprises silicon, magnesium, zinc and calcium. The content of minerals and trace elements in the preferred embodiments is preferably 1 to 80% by weight in product A and product B, more preferred 2 to 50% by weight in product A, particularly preferred 5 to 30% by weight, and 2 to 70% by weight in product B, particularly preferred 3 to 60% by weight, based on the sum of all active ingredients of product A and B, respectively. The trace elements are preferably added in the form of chromium, zinc and silicon yeasts or other organic compounds. The major elements, such as for example magnesium and calcium, are present in the form of inorganic salt compounds or organic compounds. For instance, magnesium or calcium are added as magnesium or calcium carbonate.

In the two-phase preparation of the present invention all known unsaturated fatty acids can be used. Preferably, the unsaturated fatty acids contained in vegetable and animal oils are used. Polyunsaturated fatty acids from vegetal or animal sources are essential precursors of important metabolism regulators (eicosanoids and prostaglandins). In the preferred embodiments of the two-phase preparation, the unsaturated fatty acids are preferably present in product A in an amount of 5 to 90% by weight, more preferably 10 to 70% by weight and particularly preferred 30 to 60% by weight, and in product B in a preferred amount of 1 to 90% by weight, more preferably 3 to 85% by weight and particularly preferred 5 to 80% by weight, based on the sum of all active ingredients of product A and B, respectively. Examples of preferred purely vegetal sources of unsaturated fatty acids include evening primrose oil, linseed oil, olive oil and wheat germ oil and preferred animal sources include for example fish oil.

Optionally, the two-phase preparation of the present invention can comprise all known bioactive plant substances.

The bioactive plant substances used in the present invention in particular comprise carotenoids, phytosterols, saponins, polyphenols, flavonoids, terpenes, phytoestrogens, sulfides, phytic acid and dietary fiber. Of the above-mentioned bioactive plant substances, polyphenols, flavonoids and bioflavonoids and dietary fiber are most preferably used in the present invention. Bioflavonoids of natural sources such as citrus fruit flavonoids are especially preferred in the two-phase preparation. In the preferred embodiments, the bioactive plant substances can preferably be present in product A and/or product B of the two-phase preparation in an amount of 1 to 50% by weight, more preferably 1 to 30% by weight and most preferably 1 to 20% by weight, based on the sum of all active ingredients of product A and B, respectively. Preferred natural sources of the vasco-active and antioxidant bioflavonoids are carrot, broccoli and citrus extracts as well as green tea.

The two-phase preparation of the present invention can optionally contain all known amino acids and amino acid derivatives. Preferably, in the two-phase preparation mixtures of amino acids and/or amino acid derivatives are used, including the amino acids arginine, ornithine, leucine, isoleucine, valine and tryptophan and the amino acid derivatives taurine and carnitine. Further preferred are mixtures of amino acids and amino acid derivatives with vitamin B6 and coenzyme Q<sub>10</sub> to promote regeneration and anabolism during the night phase. In the preferred embodiments, the amino acids can preferably be present individually or in the form of mixtures in product A and/or product B of the two-phase preparation in an amount of 10 to 95% by weight, more preferably 50 to 90% by weight and most preferably 80 to 90% by weight, based on the sum of all active ingredients of product A and B, respectively. The amino acids and their derivatives are preferably added in pure form.

Optionally, the two-phase preparation of the present invention can comprise all known adjuvants, additives, carriers and solvents. Preferred examples thereof include milk fat, hydrogenated, partly hydrogenated and unhydrogenated soybean fat, soybean oil, walnut butter, glycerin, gelatin, sorbitol solution or dry substance, iron oxide, titanium dioxide, patent blue, quinoline yellow, ponceau and water.

The manufacture of the two-phase preparation of the present invention can be carried out in the usual manner known to the person skilled in the art by combining the active ingredients with suitable non-toxic, inert, pharmaceutically acceptable solid or liquid carriers and optionally the usual additives, adjuvants and solvents to a galenic form of administration. Methods for manufacturing galenic forms of administration, e.g. soft gelatin capsules, are for example described in H. Sucker, P. Fuchs, P. Speiser, *Pharmazeutische Technologie* [Pharmaceutical Technology], 2<sup>nd</sup> edition 1991, Georg Thieme Verlag, Stuttgart; R. Voigt, *Lehrbuch der pharmazeutischen Technologie* [Handbook of Pharmaceutical Technology], Chemie Verlag 1976; Fahrig and Hofer *Die Kapsel* [The Capsule], Wissenschaftliche Verlagsgesellschaft 1982; Remington's *Pharmaceutical Sciences*, 15<sup>th</sup> edition 1975, Mack Publishing Company, Easton, Pennsylvania.

For the two-phase preparation of the present invention all known forms of application such as capsules, tablets, dragées and solutions etc. and all known ways of application are possible. In particular, however, oral administration is preferred, the preferred forms of application being soft gelatin capsules, ampoules for drinking, small bags (small ampoules) and ready-to-drink liquids. If, for instance, the two-phase preparation of the present invention is provided in soft gelatin capsules, the dosage is preferably such that the product A, which contains the reasonable nutrients for

the day phase or active phase, is enclosed in a capsule A which should preferably be administered in the morning. The product B, which contains the reasonable nutrients for the night phase or regenerative phase, is enclosed in a capsule B which should preferably be administered in the evening. Both capsules A and B contain the total dosage for the day and the night phases of one day, respectively. It is suggested to administer per day a capsule A in the morning and a capsule B in the evening. The two-phase preparation of the present invention can be provided as a kit, comprising for example the same number of capsules A and capsules B. Further forms of application can be ampoules for drinking and bags. However, it is also possible to provide product A and product B modified in a single form of application such that the active ingredients of product A and product B are released at different times (i.e. depot form).

The present invention is explained in more detail by means of examples.

#### Example 1

This example comprises a two-phase preparation for use at differential times, which is especially tailored to the female organism, as a dietary supplement. The compositions of capsule A and capsule B given in Table I were obtained in the usual manner known to the person skilled in the art and enclosed in soft gelatin capsules by means of common methods. Additionally, the preparation may contain known additives, adjuvants and carriers, and solvents.

Table ICapsule A

10	mg	carrot extract	
200	mg	evening primrose oil	
100	mg	linseed oil	
50	mg	soy lecithin	
5	mg	beta carotene	in the form of a 30% beta carotene suspension
30	mg	vitamin E	
200	mg	vitamin C	
200	mg	zinc yeast	(10 mg Zn/g yeast)
50	mg	magnesium	
50	mg	flavonoids	from citrus fruit

Capsule B

20	mg	broccoli extract	
200	mg	evening primrose oil	
100	mg	linseed oil	
5	mg	vitamin E	
100	mg	silicon yeast	(30 mg Si/g yeast)
300	mg	calcium	
3	mg	vitamin B1	
3	mg	vitamin B2	
5	mg	vitamin B6	
8	mg	pantothenic acid	
10	mg	niacin	
5	µg	vitamin B12	
150	µg	folic acid	
100	µg	biotin	
100	mg	zinc yeast	(10 mg Zn/g yeast)
50	mg	flavonoids	from citrus fruit



### Example 2

This example comprises a two-phase preparation for use at differential times, which is especially tailored to the male organism, as a dietary supplement. The compositions of capsule A and capsule B given in Table II were obtained in the usual manner known to the person skilled in the art and enclosed in soft gelatin capsules by means of common methods. Additionally, the preparation may contain known additives, adjuvants and carriers, and solvents.

Table II

#### Capsule A

300	mg	fish oil (icosapentenoic acid)
100	mg	linseed oil
100	mg	soy lecithin
36	mg	vitamin E
200	mg	vitamin C
5	mg	beta carotene
100	mg	magnesium
100	mg	chromium yeast
50	mg	bioflavonoids

#### Capsule B

300	mg	olive oil
300	mg	wheat germ oil
100	mg	magnesium
4.5	mg	vitamin B1
5	mg	vitamin B2
5	mg	vitamin B6
8	mg	pantothenic acid
10	mg	niacin
5	µg	vitamin B12
200	mg	zinc yeast (10 mg Zn/g yeast)

100 mg green tea extract

### Example 3

This example comprises a two-phase preparation for use at differential times, which is especially tailored to the organism of athletes, as a dietary supplement. The compositions of capsule A and bag B given in Table III were obtained in the usual manner known to the person skilled in the art. Additionally, the formulation may contain known additives, adjuvants and carriers, and solvents.

Table III

#### Capsule A

200	mg	magnesium
250	mg	linseed oil
250	mg	evening primrose oil
36	mg	vitamin E
225	mg	vitamin C
6	mg	beta carotene
4.5	mg	vitamin B1
5	mg	vitamin B2
5	mg	vitamin B6
8	mg	pantothenic acid
10	mg	niacin
5	µg	vitamin B12
200	mg	zinc yeast (10 mg Zn/g yeast)
100	mg	chromium yeast
100	mg	flavonoids from citrus fruit

#### Product B (bag)

Mixture of amino acids and amino acid derivatives with vitamin B6 and coenzyme Q<sub>10</sub> to promote regeneration and

anabolism during the night phase and to promote the fat metabolism/lipolysis.

100 mg zinc yeast (10 mg Zn/g yeast)  
100 mg magnesium  
200 mg wheat germ oil  
50 mg flavonoids from citrus fruit  
2000 mg mixture of amino acids and amino acid derivatives  
with vitamin B6 and coenzyme Q<sub>10</sub>

PCT/EP96/03329  
Schlachter, Herbert, et al.  
u.Z.: A 2376 PCT

Claims

as filed on August 14, 1997

1. Use of a product A and a product B for the preparation of a two-phase food supplement preparation for use at differential times comprising the product A for the day phase and the product B for the night phase wherein the product A comprises
  - a) at least one representative of the unsaturated fatty acids and/or
  - b) at least one representative of the group of trace elements and minerals and/or
  - c) at least one representative of the group of vitamins and/or
  - d) at least one representative of the bioactive plant substances and/or
  - e) at least one amino acid and/or amino acid derivative,and  
the product B comprises
  - a) at least one representative of the unsaturated fatty acids and/or
  - b) at least one representative of the group of trace elements and minerals and/or
  - c) at least one representative of the group of vitamins and/or
  - d) at least one representative of the bioactive plant substances and/or
  - e) at least one amino acid and/or amino acid derivative,with the proviso that product A and product B differ from each other in their quantitative and/or material composition.

2. Use according to claim 1 comprising  
a product A and a product B which comprise,  
independently of each other, the components a), b), and  
c).
3. Use according to claim 1 or 2, wherein the product A  
and/or the product B additionally comprise(s) soy  
lecithin.
4. Use according to claim 2 or 3, wherein the product A  
and/or the product B additionally comprise(s) at least  
one representative of the group of bioactive plant  
substances.
5. Use according to claims 2 to 4, wherein the product A  
and/or the product B additionally comprise(s) an amino  
acid and/or amino acid derivative or a mixture of amino  
acids and/or amino acid derivatives.
6. Use according to claim 2, comprising a product A which  
comprises the component a) in an amount of from 5 to  
90% by weight, the component b) in an amount of from 1  
to 80% by weight and the component c) in an amount of  
from 1 to 70% by weight, and  
  
a product B which comprises the component a) in an  
amount of from 1 to 90% by weight, the component b) in  
an amount of from 1 to 80% by weight and the component  
c) in an amount of from 0.5 to 50% by weight, based on  
the sum of all active ingredients of product A and B,  
respectively.
7. Use according to claim 2, comprising a product A which  
comprises the component a) in an amount of from 30 to  
60% by weight, the component b) in an amount of from 5  
to 30% by weight and the component c) in an amount of  
from 20 to 45% by weight, and

a product B which comprises the component a) in an amount of from 5 to 80% by weight, the component b) in an amount of from 3 to 60% by weight and the component c) in an amount of from 1 to 35% by weight, based on the sum of all active ingredients of product A and B, respectively.

8. Use according to claim 1 or 2, wherein the product A and/or the product B additionally comprise(s) 1 to 40% by weight of soy lecithin, based on the sum of all active ingredients of product A and B, respectively.
9. Use according to claim 1 or 2, wherein the product A and/or the product B additionally comprise(s) 1 to 50% by weight of at least one representative of the group of bioactive plant substances, based on the sum of all active ingredients of product A and B, respectively.
10. Use according to claim 1 or 2, wherein the unsaturated fatty acids of the product A and the product B are unsaturated fatty acids contained in vegetal and animal oils.
11. Use according to claim 1 or 2, wherein the trace elements and minerals of the product A and the product B are zinc, magnesium, calcium, silicon and chromium.
12. Use according to claim 1 or 2, wherein the vitamins of the product A and the product B are the vitamins B1, B2, B6, B12, C, E, pantothenic acid, niacin, folic acid and biotin.
13. Use according to claim 1, 4 or 9, wherein the bioactive plant substances of the product A and/or the product B are the bioflavonoids contained in carrot, broccoli and citrus extracts and green tea.

14. Use according to claim 1 or 5, wherein the product A and/or the product B contain arginine, ornithine, leucine, isoleucine, valine and tryptophan as amino acids and/or taurine and carnitine as amino acid derivatives, either individually or as a mixture of amino acids and/or amino acid derivatives.
15. Use of a product A and a product B as defined in claims 1 to 14 for the preparation of a two-phase drug preparation for use at differential times comprising the product A for the day phase and the product B for the night phase.
16. Kit containing a two-phase food supplement preparation comprising a product A and a product B as defined in claims 1 to 14.
17. Kit containing a two-phase drug preparation comprising a product A and a product B as defined in claims 1 to 14.